Columbia University College of Physicians and Surgeous

690 WEST 168TH STREET

NEW YORK 32, N. Y.

DEPARTMENT OF BIOCHEMISTRY

April 29, 1948

Dr. Joshua Lederberg
The University of Wisconsin
College of Agriculture
Department of Genetics
Madison 6, Wisconsin

Dear Joshua:

I should have written to you before not only to congratulate you upon the position you have but also to let you know the results we obtained in our experiments with p-fluoro phenylalanine.

After finding that this compound was relatively nontoxic, we fed it. We, however, were not able to do fluorine analysis on the proteins of the rat. I am still working on a meso fluorine method. To circumvent this difficulty, I prepared a deuterio p-fluoro phenylalanine. The deuterium was in the ring. On feeding this to rats there resulted a liver protein containing deuterium. On oxidation of the protein with chromic-sulfuric acids, the phenylalanine is oxidized to benzoic acid. Presumably the p-fluoro compound goes to p-fluorobenzoic acid. The benzoic acid fraction was isolated and contained all the deuterium originally present in the protein. This could be taken as proof that the p-fluoro phenylalanine was incorporated into the protein except that I now wonder if my starting material, p-fluoro toluene were contaminated with toluene. If this were the case, my final deuterio fluoro phenylalanine would be contaminated with deuterio phenylalanine. I don't believe this was the case. I am now synthesizing p-fluoro uluene which will certainly be free of toluene. As soon as I obtain some fluoro phenylalanine, I shall send you some.

I shall send you the reprint you asked for,

Best regards,

D. Rittenberg